

## Reagents Provided

**AEVD-AFC Substrate:** 500  $\mu$ L of 1 mM AEVD substrate peptide conjugated to 7-amino-4-trifluoromethyl coumarin (AFC)

**Cell Lysis Buffer:** 100 mL of Cell Lysis Buffer

**Reaction Buffer:** 4 x 2.0 mL vials of 2X Reaction Buffer

**DTT:** 400  $\mu$ L of a 1 M solution of dithiothreitol

Upon arrival, store the entire kit at  $-20^{\circ}$  C. After the first use, only the AFC-substrate and the DTT solution need to be returned to  $-20^{\circ}$  C storage. The remaining reagents (buffers) at may be stored at  $4^{\circ}$  C.

## References

1. Kerr, J.F.R. *et al.* (1972) Br. J. Cancer. **26**:239.
2. Ameisen, J.C. *et al.* (1995) Trends in Cell Biol. **5**:27.
3. Thompson, C.B. (1995) Science **267**:1456.
4. Vermes, I. *et al.* (1995) J. Immunol. Meth. **184**:39.
5. Darzynkiewicz, Z. *et al.* (1992) Cytometry **13**:795.
6. Fernandes-Alnemri, T. *et al.* (1996) Proc. Natl. Acad. Sci. USA **93**:7464.
7. Vincenz, C, and Dixit, V.M. (1997) J. Biol. Chem. **272**:6578.
8. Ng *et al.*, (1999) J. Biol. Chem. **274**:10301.

## Intended Use

To determine the increased enzymatic activity of the caspase-10 class of proteases in apoptotic cells by fluorometric reaction.

## Background Information

Apoptosis was originally described as a mechanism of controlled or physiological cell death (1). It is associated with the regulation of cellular homeostasis in organs and the elimination of damaged cells or of cells with deleterious reactivities from the host. Apoptosis is very common in tissues with intense hematopoietic activity (*e.g.* bone marrow and thymus) and in organs with high proliferative activity. Additionally, apoptosis has been implicated in the progression of a number of pathological conditions, including AIDS, cancer and autoimmune diseases (2, 3).

Apoptosis is characterized by a variety of cellular changes including loss of membrane phospholipid asymmetry (4), chromatin condensation, mitochondrial swelling and DNA cleavage (5). The end result of these changes is a form of cell death that avoids the normal inflammatory response associated with necrosis.

Caspase-10 is an intracellular cysteine protease that exists as a proenzyme activated during the cascade of events associated with apoptosis. Four isoforms of caspase-10 have been described: caspase-10/a also known as Mch-4 (6), caspase-10/b also known as FLICE2 (7), caspase-10/c, and caspase-10/d (8). Caspase-10 is a group III caspase highly homologous to caspase-8 (6,7). Caspase-10 contains two death effector domains and can be recruited along with caspase-8 into apoptosis signaling complexes associated with TNFR-like death receptors including Fas (6,7).

## Principle of the Test

Cells that are suspected or have been induced to undergo apoptosis are first lysed to collect their intracellular contents. The cell lysate can then be tested for protease activity by the addition of a caspase-specific peptide that is conjugated to the fluorescent reporter molecule 7-amino-4-trifluoromethyl coumarin (AFC). The cleavage of the peptide by the caspase releases the fluorochrome that when excited by light at 400 nm wavelength emits fluorescence at 505 nm. The level of caspase enzymatic activity in the cell lysate is directly proportional to the fluorescence signal detected with a fluorometer or a fluorescent microplate reader.

*Continued on next page.....*

FOR RESEARCH USE ONLY.

NOT FOR USE IN DIAGNOSTIC OR THERAPEUTIC PROCEDURES.

## ***Sample Preparation***

- 1) Cells that have been induced to undergo apoptosis are collected by centrifugation in a conical tube at 250 x g for 10 minutes (note: we recommend counting the cells before pelleting them). The supernatant is gently removed and discarded while the cell pellet is lysed by the addition of the Cell Lysis Buffer. The amount of Cell Lysis Buffer to be added to the pellet is determined by the number of cells present (this can be estimated from the number of cells initially cultured). Add 25  $\mu$ L of cold Cell Lysis Buffer per  $1 \times 10^6$  cells.
- 2) The cell lysate is incubated on ice for 10 minutes. This should yield a cell lysate with an approximate protein concentration of 2-4 mg/mL. (The protein content of the cell lysate can be estimated using a protein determination assay that is compatible with detergents present in the Cell Lysis Buffer, e.g. BCA Protein Assay, Pierce Chemical Co., catalog# 23225).
- 3) The enzymatic reaction for caspase activity is best carried out in a 96 well flat bottom microplate that can be read with a microplate reader equipped with fluorescence detection capabilities.
- 4) Each reaction requires 50  $\mu$ L of cell lysate (*i.e.* derived from  $2 \times 10^6$  or 100 - 200  $\mu$ g of total protein). If larger volumes of cell lysate are necessary to meet the above requirements, the total reaction volume may be scaled up (*i.e.* the volume of each reagent added should be proportionally increased).
- 5) Each reaction also requires 50  $\mu$ L of 2X Reaction Buffer. Prior to using the 2X Reaction Buffer, add 10  $\mu$ L of fresh DTT stock per 1 mL of 2X Reaction Buffer.
- 6) To each reaction well, add 5  $\mu$ L of caspase-10 fluorogenic substrate (AEVD-AFC).
- 7) Incubate the plate at 37°C for 1 - 2 hours.
- 8) Read the plate on a fluorescent microplate reader using filters that allow light excitation at 400 nm wavelength and can collect emitted light at 505 nm wavelength.
- 9) Additional controls that should be included in this assay are a) no cell lysate and b) no substrate. The total reaction volume must be kept constant and therefore distilled water can be used to replace the volume normally occupied by either the cell lysate or the substrate reagent.
- 10) For comparative analysis, the above assay should be repeated with non-induced cells.

This protocol may require modification, depending upon final utilization.

The results are best expressed as fold increase in caspase activity of apoptotic cells over that of non-induced cells. If the background controls (reactions where no cell lysate is added or where no AEVD-AFC substrate is added) give a substantial reading, it is recommended that these values be subtracted from the experimental results prior to calculating the fold increase.